

Serial No.:  
Exp. Date:

**Indications:** Recommended for the vaccination of healthy, susceptible enterotoxemia and tetanus caused by the toxins of *Clostridium perfringens* Types C and D and *Clostridium tetani*. Although *Cl. perfringens* Type B is not a significant problem in the U.S.A., immunity may be provided against the beta and epsilon toxins elaborated by *Cl. perfringens* Type B. This immunity is derived from the combination of Type C (beta) and Type D (epsilon) fractions.

**Composition:** Prepared from cultures of the organisms listed. Alum precipitated.

**Dosage:**

**Cattle:** Using aseptic technique, inject 5 mL subcutaneously. Repeat in 21 to 28 days and once annually.

**Clostridium Perfringens  
Types C & D-  
Tetanus Toxoid**  
**Bar Vac® CD/T**

250 mL/50 Cattle Doses or  
125 Sheep/Goat Doses

**Sheep and Goats:** Using aseptic technique, inject 2 mL subcutaneously. Repeat in 21 to 28 days and once annually.

**Precautions:** Store out of direct sunlight at 35-45°F (2-7°C). Avoid freezing. Shake well before using. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur.

**Antidote:** Administer epinephrine.

**Boehringer Ingelheim Vetmedica, Inc.**  
St. Joseph, Missouri 64506 U.S.A.  
U.S. Veterinary License No. 124

Code 120-321; 20302-04



**Boehringer  
Ingelheim**

**Boehringer Ingelheim Vetmedica, Inc.**

2621 North Belt Highway • St. Joseph, MO 64506-2002

Information 800-325-9167 • Sales Service 800-325-9167 • Fax 816-236-2731

[www.bi-vetmedica.com](http://www.bi-vetmedica.com)