

ZOETIS INC.

333 PORTAGE STREET, KALAMAZOO, MI, 49007

Telephone: 269-833-4000
Customer Service: 800-733-5500 and 800-793-0596
Veterinary Medical Investigations & Product Support: 800-366-5288
Technical Services (USA): 800-366-5288
Website: www.zoetis.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

QUEST[®] 2% EQUINE ORAL GEL

Zoetis

moxidectin

NADA 141-087, Approved by FDA

Contains 20 mg moxidectin/mL (2.0% w/v)

DEWORMER & BOTICIDE

FOR ORAL USE IN HORSES AND PONIES SIX MONTHS OF AGE AND OLDER

INDICATIONS

QUEST (moxidectin) 2% Equine Oral Gel when administered at the recommended dose level of 0.4 mg moxidectin/kg (2.2 lb) body weight is effective in the treatment and control of the following stages of gastrointestinal parasites in horses and ponies:

Large Strongyles:

Strongylus vulgaris - (adults and L₄/L₅ arterial stages); *Strongylus edentatus* - (adults and tissue stages); *Triodontophorus brevicauda* - (adults); *Triodontophorus serratus* - (adults)

Small Strongyles (adults): *Cyathostomum* spp., including *Cyathostomum catinatum*, *Cyathostomum pateratum*; *Cylicostephanus* spp., including *Cylicostephanus calicatus*, *Cylicostephanus goldi*, *Cylicostephanus longibursatus*, *Cylicostephanus minutus*; *Cylicocyclus* spp., including *Cylicocyclus insigne*, *Cylicocyclus leptostomum*, *Cylicocyclus nassatus*; *Cylicocyclus radiatus*; *Coronocyclus* spp., including *Coronocyclus coronatus*, *Coronocyclus labiatus*, *Coronocyclus labratus*; *Gyalocephalus capitatus*; *Petrovinema poculatus*

Small Strongyles:

Undifferentiated luminal larvae

Encysted cyathostomes

Late L₃ and L₄ mucosal cyathostome larvae

Ascarids

Parascaris equorum - (adults and L₄ larval stages)

Pinworms

Oxyuris equi - (adults and L₄ larval stages)

Hairworms

Trichostrongylus axei - (adults)

Large-mouth stomach worms

Habronema muscae - (adults)

Horse stomach bots

Gasterophilus intestinalis - (2nd and 3rd instars); *Gasterophilus nasalis* - (3rd instars)

One administration of the recommended dose rate of QUEST (moxidectin) 2% Equine Oral Gel also suppresses strongyle egg production through 84 days.

QUEST is indicated for use in horses and ponies, including breeding mares and stallions, and foals six months of age and older.

STRATEGIC PROTECTION PROGRAMS

Mature	450	(204)	2	900	(409)	1	1300	(590)	*
--------	-----	-------	---	-----	-------	---	------	-------	---

* A full syringe plus a portion of a second syringe is required to treat horses weighing more than 1150 lb.

ANIMAL SAFETY

QUEST (moxidectin) 2% Equine Oral Gel can be safely administered at the recommended dose of 0.4 mg moxidectin/kg body weight to horses and ponies of all breeds at least 6 months of age or older. Transient depression, ataxia and recumbency may be seen when very young or debilitated animals are treated. In these instances, supportive care may be advisable. Reproductive safety studies demonstrate a wide margin of safety when the product is used in the treatment of estrual and pregnant mares and breeding stallions.

To report adverse drug reactions or to obtain a copy of the Material Safety Data Sheet (MSDS) call (800)477-1365.

ENVIRONMENTAL SAFETY

Care should be taken to avoid the release of significant volumes of moxidectin into either ground or free-running water since moxidectin may be injurious to aquatic life. SURE-DIAL[®] syringes and their contents should be disposed of in an approved landfill or by incineration.

PRECAUTIONS:

QUEST 2% Equine Oral Gel has been formulated specifically for use in horses and ponies only. **This product should not be used in other animal species as severe adverse reactions, including fatalities in dogs, may result.**

WARNINGS

Extreme caution should be used when administering the product to foals, young and miniature horses, as overdosage may result in serious adverse reactions. Do not use in sick, debilitated, or underweight animals. Not for horses or ponies intended for human consumption.

HUMAN WARNINGS

Not for use in humans. Keep this and all drugs out of the reach of children. Do not ingest. If swallowed, induce vomiting. Wash hands and contaminated skin with soap and water. If accidental contact with eyes occurs, flush repeatedly with water. If irritation or any other symptom attributable to exposure to this product persists, consult your physician.

HOW SUPPLIED

QUEST 2% Equine Oral Gel is available in one syringe applicator size. Each SURE-DIAL[®] syringe contains 0.4 oz (11.3 g) of QUEST 2% Equine Oral Gel which is sufficient to treat a single horse weighing up to 1150 lb, or two or more lighter animals with a combined body weight of up to 1150 lb. Available in 20-pack displays and 50-pack bulk.

NDC 0856-7441-03 - 0.4 oz (11.3 g) syringe - 20 mg moxidectin per mL

Store at or below 77°F (25°C). Avoid freezing. If frozen, thaw completely before use. Store partially-used syringes with the cap tightly secured.

U.S. Patent No. 4,916,154

© 2005 Fort Dodge Animal Health. All Rights Reserved.

Fort Dodge Animal Health, Fort Dodge, Iowa 50501 USA

03476

Revised December 2006

7440N

NAC No.: 1003178.8