NDC 30798-237-10

durvet

Duramycin 72-200



Oxytetracycline Injection (200 mg/mL) Antibiotic

ANADA 200-306, APPROVED BY FDA

Each mL contains 200 mg of oxytetracycline base as amphoteric oxytetracycline.
For the treatment of disease in beef cattle; dairy cattle; calves, including preruminating (veal) calves; and swine.

For animal use only

Net Contents: 100 mL

Duramycin 72-200 is a sterile, preconstituted solution of the broad-spectrum antibiotic oxytetracycline.

CAUTION:
When administered to cattle, muscle discoloration may necessitate trimming of the injection site(s) and surrounding tissues during the dressing procedure.

WARNING:
Discontinue treatment at least 28 days prior to slaughter of cattle and swine. Milk taken from animals during treatment and for 96 hours after the last treatment must not be used for food.

PRECAUTIONS:
Exceeding the highest recommended level of drug per lb of body weight per day, administering more than the recommended number of treatments, and/or exceeding 10 mL intramuscularly or subcutaneously per injection site in adult beef cattle and dairy cattle, and 5 mL intramuscularly per injection site in adult swine, may result in antibiotic residues beyond the withdrawal period.

Refer to Package Insert for Complete Directions.

Storage:
Store at room temperature 15° - 30°C (59° - 86°F).
Keep from freezing.

Restricted Drug(s) (California),

Not for Human Use,

Use Only as Directed.

Manufactured for:
DURVET, INC.
Blue Springs, Missouri 64014
www.durvet.com

Made in the U.K.

NDC 30798-237-10

durvet

Duramycin 72-200



Oxytetracycline Injection (200 mg/mL) Antibiotic

ANADA 200-306, APPROVED BY FDA

Each mL contains 200 mg of oxytetracycline base as amphoteric oxytetracycline.
For the treatment of disease in beef cattle; dairy cattle; calves, including preruminating (veal) calves; and swine.

For animal use only

Net Contents: 100 mL

CATTLE DOSAGE GUIDE
At the first signs of pneumonia or pinkeye,* administer a single dose of Duramycin 72-200 by deep intramuscular injection, or subcutaneously, according to the following weight categories. **

Animal Weight (lb)	Number of mL or cc
100	4.5
200	9.0
300	13.5
400	18.0
500	22.5
600	27.0
700	31.5
800	36.0
900	40.5
1000	45.0
1100	49.5
1200	54.0

* See package insert for dosing instructions for other indicated diseases and full product information.
** Do not administer more than 10 mL at any one injection site (1-2 mL per site in small calves).
Discontinue treatment at least 28 days prior to slaughter.

SWINE DOSAGE GUIDE
At the first signs of pneumonia, * administer Duramycin 72-200 by deep intramuscular injection, according to the following weight categories. **

Animal Weight (lb)	Number of mL or cc
10	0.5
25	1.1
50	2.3
75	3.4
100	4.5
125	5.6
150	6.8
175	7.9
200	9.0
225	10.1
250	11.3
275	12.4
300	13.5
325	14.6

* See package insert for dosing instructions for other indicated diseases and full product information.
** Do not administer more than 5 mL at any one injection site.
Discontinue treatment at least 28 days prior to slaughter.

Batch No:

Exp:

PMS 109 C
PMS 185C
PMS BLACK C

000000CO1
Iss: 12/07