

CALF181

# Augusta Startem Right 18% Calf

Textured

Formulated for beef and dairy calves.

## MEDICATED

For slaughter, stocker and feeder cattle, and calves, including veal calves)for prevention of coccidiosis due to E. bovis and E. zuernii.  
Prevents the development of house flies, stable flies, face flies, and horn flies in the manure of treated cattle.

### ACTIVE DRUG INGREDIENT

Monensin (as Monensin Sodium) ..... 35 g/ton

### ACTIVE INGREDIENT

Diflubenzuron ..... 3.05 mg/lb

### Guaranteed Analysis

Crude Protein		(Min)	18.0 %
Crude Fat		(Min)	3.0 %
Crude Fiber		(Max)	6.5 %
Calcium	(Min)	0.9	(Max) 1.6 %
Phosphorus		(Min)	0.3 %
Salt	(Min)	0.4	(Max) 1.1 %
Potassium		(Min)	0.8 %
Vitamin A		(Min)	5.5 KIU/lb

### List of Ingredients

Grain Products, Plant Protein Products, Roughage Products, Brewer's Dried Yeast and Dried Saccharomyces Cerevisiae Fermentation Solubles, Vitamin A Supplement, Vitamin D3 Supplement, Vitamin E Supplement, Active Dried Yeast (Saccharomyces Cerevisiae), Dried Trichoderma Longibrachiatum Fermentation Product, Niacin, Vitamin B-12 Supplement, Riboflavin-5-Phosphate, D-Calcium Pantothenate, Choline Chloride, Biotin, Thiamine Mononitrate, Pyridoxine Hydrochloride, Folic Acid, Calcium Aluminosilicate, Sodium Aluminosilicate, Diatomaceous earth, Calcium Carbonate, Rice Hulls, Mineral Oil, Propionic Acid, Amorphous Silicon Dioxide, Verxite, Hydrated Sodium Calcium Aluminosilicate, Water, Ammonium Hydroxide, Active Dry Yeast, Yeast Culture, Dried Enterococcus Faecium Fermentation Product, Dried Lactobacillus Acidophilus Fermentation Product, Dried Lactobacillus Plantarum Fermentation Product, Dried Lactobacillus Brevis Fermentation Product, Dried Bacillus Lentus Fermentation Product, Dried Bacillus Subtilis Fermentation Product, Dried Aspergillus Oryzae Fermentation Extract, Dried Trichoderma Longibrachiatum Fermentation Extract, Salt of Glutamic Acid, Dried Grain Products, Dried Molasses Products, Mineral Oil., Cane Molasses, Vegetable Fat Product (Feed Grade), Propylene Glycol, Lecithin, Phosphoric Acid, Sulfuric Acid, Caramel Flavoring, Acetic Acid, and Benzoic Acid (as Preservatives), Processed Grain By-Products, Salt, Molasses Products, Magnesium Sulfate, Potassium Sulfate, Magnesium Oxide, Zinc Sulfate, Ethylenediamine Dihydriodide, Copper Sulfate, Ferrous Sulfate, Cobalt Carbonate, Sodium Selenite.

### Feeding Directions

Feed continuously with good quality forage and fresh, clean water.

**CAUTION:** Keep out of reach of children. Discard empty container according to local regulations. Never reuse empty container.

**CAUTION:** Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Do not exceed the levels of Monensin recommended by the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

**WARNING:** A withdrawal time has not been established for preruminating calves. Do not use in calves to be processed for veal.

**LIMITATIONS:** For calves, excluding veal calves, feed at a rate of 0.14 to 1.0 milligrams Monensin per pound of body weight per day, depending upon the severity of challenge, up to maximum of 200 milligrams per head per day.

This product contains added copper. Do not feed to sheep.

Follow label directions.

Manufactured By:

Augusta Cooperative Farm Bureau, Inc.

Staunton, Virginia 24401

NET WEIGHT 50 LBS. (22.68 kg)

## MEDICATED